

MODULATION OF ENZYME ACTIVITY

Effectors

← Inhibitor:
decreases
reaction rate

v_i

Degree of inhibition:


$$\epsilon_i = \frac{v_0 - v_i}{v_0}$$

→ Activator:
increases
reaction rate

v_a

Degree of activation:

$$\epsilon_a = \frac{v_a - v_0}{v_0}$$



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

INHIBITION

REVERSIBLE

$$E + S \rightleftharpoons ES \longrightarrow E + P$$


↓↑
EI


IRREVERSIBLE

$$E + S \xrightleftharpoons{K_s} ES \xrightarrow{k_2} E + P$$

↓
EI

distinction:






ny Tanszék

Competitive inhibition

Competition between S and I for the active sites of the enzyme, or mutual exclusion



- I may be an:
- substrate analogue
 - alternative substrate
 - product

MODEL 1.: Classical competitive inhibition:

I competes with S for occupation of the same active site




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

3

COMPETITIVE INHIBITION

MODEL 2.: steric hindrance A



Binding of I to another site sterically hinders S in binding to the active site of enzyme.




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

4

COMPETITIVE INHIBITION

MODEL 3.: steric hindrance B



An analog part of S and I compete for a common binding site.




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

5

COMPETITIVE INHIBITION

MODEL 4.: overlapping



Sites 1 and 3 can bind I, 2 and 4 can bind S, but both exclude each other.



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék


6

COMPETITIVE INHIBITION

MODEL 5.:

Binding of **I** changes the conformation of the enzyme which prevents binding of **S** to the active centre.

End product inhibition (feed back inhibition) is typical example of this case.




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

7

Kinetics of competitive inhibition

Basic equations for competitive inhibition:



$$K_s = \frac{E \cdot S}{(ES)}$$

$$K_i = \frac{E \cdot I}{(EI)}$$

- if $k_{app} > 0$ than **I** is an alternative substrate
- if $k_{app} = 0$ than **I** is a „dead end” competitive inhibitor




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

8



Kinetics of competitive inhibition

Alternative substrate: the enzyme is able to transform the structural analogous molecule, too. → an *alternative product* is formed.



Enzymes with group and region specificity have numerous alternative substrates

Example: the enzymes of liver: alcohol dehydrogenase, aldehyde dehydrogenase:




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

9

Kinetics of competitive inhibition

Repeat the deduction:



$$K_s = \frac{E \cdot S}{(ES)}$$

$$K_i = \frac{E \cdot I}{(EI)}$$

product formation
rate:

$$V = \frac{dP}{dt} = k_2 (ES)$$

Mass balance of enzyme:

$$E_0 = E + (ES) + (EI)$$



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

10

Kinetics of competitive inhibition

Divide the two equation:

$$\frac{V}{E_o} = \frac{k_2(ES)}{E + (ES) + (EI)}$$

Substitute:

$$K_s = \frac{E \cdot S}{(ES)} \quad K_i = \frac{E \cdot I}{(EI)}$$

$$\frac{V}{E_o} = \frac{k_2 E \frac{S}{K_s}}{E + E \frac{S}{K_s} + E \frac{I}{K_i}} \quad \rightarrow \quad V = \frac{k_2 E_o \frac{S}{K_s}}{1 + \frac{S}{K_s} + \frac{I}{K_i}}$$

$$V_{\max} = k_2 E_o$$



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

11

Kinetics of competitive inhibition

Simplified forms of reaction rate:

$$\frac{V}{V_{\max}} = \frac{S}{K_s \left(1 + \frac{I}{K_i} \right) + S}$$

or:

$$V = V_{\max} \frac{S}{K_s \left(1 + \frac{I}{K_i} \right) + S}$$


or:

$$v_i = \frac{v_{\max}(S)}{K_s \left[\frac{K_i + (I)}{K_i} \right] + (S)}$$



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

12



Competitive inhibition

Alternative substrates: for hexokinase: glucose, fructose

S-analogons: drugs:

$$\begin{array}{c}
 \text{H} \quad \text{H} \\
 | \quad | \\
 \text{HC} - \text{C} - \text{NH}_2 \\
 | \\
 \text{C} \\
 / \quad \backslash \\
 \text{HO} \quad \text{O}
 \end{array}$$

L-alanin

$$\begin{array}{c}
 \text{H} \quad \text{H} \\
 | \quad | \\
 \text{HC} - \text{C} - \text{NH}_2 \\
 | \quad | \\
 \text{O} \quad \text{C} \\
 \backslash \quad / \\
 \quad \text{N} \\
 | \\
 \text{H}
 \end{array}$$

cikloszerin

BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

Competitive inhibition

Effect of sulfamides (antimicrobial drugs): substrate analogon act as competitive inhibitor.

p-amino benzoic acid

↓ **PABA CONVERTED TO FOLIC ACID**

Folic acid **REQUIRED BY BACTERIA**

Sulfameth oxazole **ANALOGUE OF PABA BLOCKS SYNTHESIS OF F.A.**

BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

Analogous inhibitions

competitive inhibition:

$$V = V_{\max} \frac{S}{K_s \left(1 + \frac{I}{K_i} \right) + S}$$

product inhibition:


$$V = V_{\max} \frac{S}{K_s \left(1 + \frac{P}{K_p} \right) + S}$$

alternative or competing substrates

$$V_1 = V_{1\max} \frac{S_1}{K_{S1} \left(1 + \frac{S_2}{K_{S2}} \right) + S_1}$$

$$V_2 = V_{2\max} \frac{S_2}{K_{S2} \left(1 + \frac{S_1}{K_{S1}} \right) + S_2}$$

BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék



Noncompetitive inhibition

Inhibitor binds to an other active site of the enzyme and does not affect the binding of the substrate – does not change the affinity of the enzyme to the substrate.

It exists only when rapid equilibrium can be supposed, $K_s = K_m$.

Equations of noncompetitive inhibition:

$$E + S \xrightleftharpoons{K_s} ES \xrightarrow{k_p} E + P$$

$$EI + S \xrightleftharpoons{K_s} ESI$$

$$K_s = \frac{E \cdot S}{ES} = \frac{EI \cdot S}{ESI} \quad \text{és} \quad K_i = \frac{E \cdot I}{EI} = \frac{ES \cdot I}{ESI}$$

$$V = k_p(ES)$$

$$\frac{V}{V_{max}} = \frac{ES}{E + ES + EI + ESI}$$

BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

Noncompetitive inhibition

$$\frac{V}{V_{\max}} = \frac{\frac{S}{K_s}}{1 + \frac{S}{K_s} + \frac{I}{K_i} + \frac{S \cdot I}{K_s K_i}}$$

or

$$\frac{V}{V_{\max}} = \frac{S}{K_s \left(1 + \frac{I}{K_i}\right) + S \left(1 + \frac{I}{K_i}\right)}$$

or

$$V = V_{\max} \frac{1}{\left(1 + \frac{I}{K_i}\right)} \frac{S}{K_s + S}$$


$$\frac{V}{V_{\max}} = \frac{ES}{E + ES + EI + ESI}$$

Inhibitor changes the value of the apparent V_{\max} , but does not change the values of K_s (or K_m).

$$V = V_{\max i} \frac{S}{K_s + S} \quad \text{where } V_{\max i} = V_{\max} \frac{1}{1 + \frac{I}{K_i}}$$

BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

19



Noncompetitive inhibition

Examples:

H⁺ ions' effect on chymotripsine. Here a proton acceptor site exists in the active centre, which can be inhibited by increasing H⁺-ion concentration. (L-B plot shows clear noncompetitive inhibition, (but do not forget the complex effect of the pH on enzymes).

Heavy metal molecules(-SH reagensek), or cyanides.
Often these effects are irreversible.




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

21

Noncompetitive inhibition

Surface of slices apple gets brown in air: o-diphenol oxidase enzyme catalyses the catechol → o-quinone reaction



this and other reaction products give the brown color

competitive inhibitor of o-diphenol oxidase is para-hydroxybenzoic acid (PHBA), a structural analog.





BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

22

Noncompetitive inhibition

competitive inhibitor of o-diphenol oxidase is para-hydroxybenzoic acid (PHBA), a structural analogon

noncompetitive inhibitor is: phenylthiourea, bound to copper ion what is necessary to enzyme activity.




BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

23

Uncompetitive inhibition

Fixed order: the inhibitor must join second, after the substrate




24

Uncompetitive inhibition


$$V = V_{\max} \frac{1}{1 + \frac{I}{K_i}} \cdot \frac{S}{\left(1 + \frac{I}{K_i}\right) K_m + S}$$

$$\frac{1}{V} = \frac{K_m}{V_{\max}} \frac{1}{S} + \frac{1}{V_{\max}} \left(1 + \frac{I}{K_i}\right)$$



Linear mixed type inhibition

Mechanism of linear mixed type inhibition resembles to non-competitive inhibition but presence of I modifies the enzyme affinity to substrate.



Linear mixed type inhibition

Expressing the change of two kinetic parameters:

$$V = V_{\max} \frac{1}{\left(1 + \frac{I}{\alpha K_I}\right)} \cdot \frac{S}{K_s \cdot \frac{\left(1 + \frac{I}{K_I}\right)}{\left(1 + \frac{I}{\alpha K_I}\right)} + S}$$

$$V_{\max i} = V_{\max} \frac{1}{\left(1 + \frac{I}{\alpha K_I}\right)}$$

$$K_{s i} = K_s \cdot \frac{\left(1 + \frac{I}{K_I}\right)}{\left(1 + \frac{I}{\alpha K_I}\right)}$$



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

27

competitive

$$V = V_{\max} \frac{S}{K_s \left(1 + \frac{I}{K_I}\right) + S}$$

noncompetitive

$$V = V_{\max} \frac{1}{\left(1 + \frac{I}{K_I}\right)} \cdot \frac{S}{K_s + S}$$

uncompetitive

$$V = V_{\max} \frac{S}{K_s + S \left(1 + \frac{I}{K_I}\right)}$$

mixed

$$V = V_{\max} \frac{S}{K_s \left(1 + \frac{I}{K_I}\right) + S \left(1 + \frac{I}{\alpha K_I}\right)}$$

$$V = V_{\max} \frac{1}{\left(1 + \frac{I}{\alpha K_I}\right)} \cdot \frac{S}{K_s \cdot \frac{\left(1 + \frac{I}{K_I}\right)}{\left(1 + \frac{I}{\alpha K_I}\right)} + S}$$

$$V = V_{\max} \frac{1}{1 + \frac{I}{K_I}} \cdot \frac{S}{\frac{K_m}{\left(1 + \frac{I}{K_I}\right)} + S}$$



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

28

Summary of the inhibition types

S and I mutually exclude each other from the enzyme
COMPETITIVE

S and I bind to the enzyme independently on each other
NONCOMPETITIVE

I binds only after S
UNCOMPETITIVE

Like former but I modifies the affinity of the enzyme
MIXED TYPE



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék


29

Substrate inhibition

The substrate binds to two or more sites.

If the S concentration is high, it can occur that two S bind to one and the other binding site forming inactive complex.

(also reversible inhibition).



BME Alkalmazott Biotechnológia és Élelmiszertudomány Tanszék

30

