

6. OTHER EFFECTS ON ENZYME ACTIVITY

- Ionic strength
- pH
- TEMPERATURE
- Shear
- Pressure (hydrostatic)
- Surface tension
- Chemicals (alcohol, urea, H_2O_2 ...)
- Light, sonication, ionising radiations


Reversible changes
Irreversible



BME Alkalmazott Biotechnológia és Élelmisztudomány Tanszék

1

Effect of pH



$$Y^- = \frac{1}{1 + H^+ / K_1 + K_2 / H^+}$$

$$H^+_{\text{optimum}} = \sqrt{K_1 K_2}$$

$$(pH)_{\text{optimum}} = \frac{1}{2} (pK_1 + pK_2)$$

$$V_{\max} = k_2 E_0 Y^- = k_2 E_0 \frac{1}{1 + H^+ / K_1 + K_2 / H^+}$$



4

Active side chains

Changes in activity of proteins are caused by changes of amino acid side chains.

Acidic: $-\text{COOH}$: Asp, Glu Basic: $-\text{NH}_2$: Lys, Arg
(and terminal $-\text{COOH}$ and $-\text{NH}_2$)

amide: $-\text{CO}-\text{NH}_2$: Asn, Gln

Polar: $-\text{OH}$: Ser, Thr $-\text{SH}$: Cys, $-\text{S}-\text{CH}_3$: Met

Imidazole: His Guanidin: Arg


H-bonds: $\text{C=O} \dots \text{H-O-}$ $\text{C=O} \dots \text{H-NH-}$



BME Alkalmazott Biotechnológia és Élelmisztudomány Tanszék

2

Effect of pH




5

Effect of pH

Proteins: + and - charged side chains \leftarrow their charge depends on dissociation \leftarrow determined by pH \rightarrow it effects the active centre.

Recharge of enzyme:



Only E^- is active!

Ratio of active enzymes: $Y^- = \text{E}^- / E_0$

Michaelis-féle pH függvények:


$$Y^- = \frac{1}{1 + H^+ / K_1 + K_2 / H^+}$$



BME Alkalmazott Biotechnológia és Élelmisztudomány Tanszék

3

Effect of pH temperature



$$\frac{dE_a}{dt} = -kE_a \quad \longrightarrow \quad E_a(t) = E_{a0} e^{-kt}$$

6

Effect of pH temperature

$E_a \rightleftharpoons E_i$

$$\frac{E_i}{E_a} = K_d = \exp\left(\frac{-\Delta G_d}{RT}\right) = \exp\left(\frac{-\Delta H_d}{RT}\right) \exp\left(\frac{\Delta S_d}{R}\right)$$

If: $V_{max} = k_2(T)E_a$

$$E_0 = E_a + E_i \rightarrow E_a = \frac{E_0}{1+K_d}$$

and $k_2(T) = \beta \left(\frac{k_B T}{h}\right) e^{\Delta S^\circ / R} \cdot e^{-\Delta H_d / RT}$

$S_d = \sim 900 \text{ KJ/mol.K}$
 $H_d = 280-310 \text{ KJ/mol}$

Large: sensitively reacts on small change (one H-bond: 12,5-29,3 KJ/mol)

$$V_{max} = \frac{\alpha T e^{-E/RT}}{1 + e^{\Delta S^\circ / R} \cdot e^{-\Delta H_d / RT}}$$

$\alpha = \text{combination of } (\beta, k_B, h, E_0, \Delta S^\circ)$

K_m also depends on T !

BME Alkalmazott Biotechnológia és Élelmiszer tudomány Tanszék

Temperature optimum and stability

Temperature °C	20	30	40	50	60	70	80	90	100	110
Microbial rennet	—	—	—	—	—	—	—	—	—	—
<i>A. oryzae</i> α -amylase	—	—	—	—	—	—	—	—	—	—
<i>B. subtilis</i> α -amylase	—	—	—	—	—	—	—	—	—	—
<i>B. licheniformis</i> α -amylase	—	—	—	—	—	—	—	—	—	—
<i>A. oryzae</i> β -galactosidase	—	—	—	—	—	—	—	—	—	—

Stability optimum (pink line)
Activity optimum (red line)
(optimum set at $\geq 80\%$ of maximum)

BME Alkalmazott Biotechnológia és Élelmiszer tudomány Tanszék

